

# The Relationship Between the Level of Depression and Anxiety in Parents with Primary School Children, and Preschool Children Diagnosed with Neuro-Developmental Disorders

Lon LASKA<sup>1\*</sup>, Afërdita HABIBAJ<sup>2</sup>, Afërdita AHMETI<sup>3</sup>, Shpresë QAMILI<sup>4</sup>

<sup>1</sup>Professor, Faculty of Applied Psychology and Management Studies, College "DARDANIA", Prishtina, Kosovo;

<sup>2</sup>Faculty of General psychology, Mitrovica, Kosovo;

<sup>3</sup>Department of Psychology, AAB college, Prishtina, Kosovo;

<sup>4</sup>Prof. Assoc. Dr. Shpresë Qamili, Faculty of Applied Sciences, Public University " Kadri Zeka", Gjilan, Kosovo

## ABSTRACT

This study aimed to evaluate the relationship between depression and anxiety in parents with children diagnosed with neurodevelopmental disorders. The main purpose of this study is to measure the level of depression and anxiety in parents whose primary school children and preschool children have been diagnosed with neurodevelopmental disorders and to correlate these disorders with sociodemographic characteristics. In this study participated 300 parents whose children were diagnosed with neuro-developmental disorders. For the methodology of this study, the integration of quantitative and qualitative methods was used. In this study, two closed-type instruments were used, one instrument measures anxiety and the other depression in parents whose primary school children and preschool children have been diagnosed with neurodevelopmental disorders. The findings of this study have shown that depression has a high positive correlation with anxiety. As the level of depression in parents increases, so does the level of anxiety. Gender differences, statistically significant, between the level of depression of parents whose children have been diagnosed with neuro-developmental disorders, we see that mothers reach a higher average than fathers, also when it comes to anxiety, mothers reach a higher average than fathers.

**Keywords:** Depression, anxiety, parents, *primary school children, preschool children, neuro-developmental, disorders.*

## INTRODUCTION

In this paper, we will talk about the relationship between depression and anxiety in parents with children diagnosed with neuro-developmental disorders as well as their causes and consequences.

Considering that these disorders are very current, today each of us should be informed about depression and anxiety and their impact on our lives, autism how it appears, and why it appears (Stoep & Weiss, 2003).

One of the main reasons why I chose such a topic is that families of children with mental health problems refer to increased weight/burden, loss of work, and increasing parent-child conflicts. (Brannan & Heflinger, 2006).

From other studies, we see that depression is a mental state of the individual, which appears with negative changes and with a feeling of fear, internal discomfort, as well as obstacles in thinking.

These changes are manifested in the reduction of the individual's interest in events, in the loss of initiative to act, in despair and in a pessimistic attitude toward the future. Depressed people have feelings of guilt, thoughts, and ideas of the sinner, which can rise to delusional states. Obstacles in sleep are expressed by difficulties during sleep, insomnia, and repeated and early awakening, so there is no freshness in the morning. The morning is often the worst time for depressed patients, as that's when they feel really bad. At lunch, their

condition improves and in the evening it becomes relatively bearable. (Stoep & Weiss, 2003).

Anxiety is defined as a condition, which can be caused as a result of interpersonal relationships. Anxiety is perceived as an undesirable emotional state, which is characterized by the subjective emotional feeling of tension, evaluation of the situation, and worry, as well as by the activation of the autonomic nervous system, which can be found in a more stable form, especially in its use in the mental health literature. Feelings of anxiety come as a result of the nature of the relationships that the individual has and creates with others.

**Corresponding Author:** lon.laska@hotmail.com

<https://orcid.org/0000-0002-0511-9786>

**How to cite this article:** LASKA L, HABIBAJ A, AHMETI A, QAMILI S (2023). The Relationship Between the Level of Depression and Anxiety in Parents with Primary School Children, and Preschool Children Diagnosed with Neuro-Developmental Disorders. Pegem Journal of Education and Instruction, Vol. 13, No. 2, 2023, 21-27

**Source of support:** Nil

**Conflict of interest:** None.

**DOI:** 10.47750/pegegog.13.02.03

**Received :** 10.10.2022

**Accepted :** 16.19.2022

**Published:** 01.03.2023

So, if these relationships are filled with tension, stress, disagreement or aggressiveness, then the individual experiences similar situations with anxiety, or will respond in the same way to new situations in his life. (Betty, 1997). This study also emphasizes the awareness and psychoeducation of parents on positive parenting strategies for raising children with neurodevelopmental disorders. Mental health problems, defined as neuro-developmental or cognitive, even associated with impairment (Sayal, 2006), affect every aspect of children’s development and functioning, including relationships with peers and family, behavior at school, and the transition to adulthood. For example, children with mental health problems are at greater risk than those without problems to drop out of school (Stoep & Weiss, 2003), at risk of becoming victims through physical or sexual abuse (Unger & Kipke, 1997), to have conflicts with parents (Verhulst & van der Ende, 1997), to be drug and alcohol users (White, Xie et al., 2001), and to have problems with justice (Cocozza & Skowrya, 2000; Huber & Wolfson, 2000), be violent or die by suicide. Families of children with mental health problems report increased weight/burden, job loss, and increased parent-child conflict (Brannan & Heflinger, 2006). The societal cost of children’s mental health problems can be calculated in the increased frequency of academic failure, youth incarceration, suicide attempts, and psychiatric hospitalizations (US Public Health Service, 2000).

The results of this study may improve the future treatment of children with neurodevelopmental disorders and help parents manage anxiety and depression. (Stoep & Weiss, 2003). The review of the literature on depression and anxiety in parents with children diagnosed with neuro-developmental disorders, with a research character, is developed according to the perspectives of different theories of depression and anxiety presented in the theoretical bases. Different studies on influencing factors are examined by comparing and contrasting them with each other. The findings and results evidenced by the reviewed literature serve as the initiative and commitment of this paper to provide valuable assistance with new findings and recommendations in this field.

## RESEARCH METHODOLOGY

In this study, we used quantitative research, as it provides several advantages compared to other research alternatives and I believe it has best achieved the purpose of the study. The main advantage of quantitative research is that it provides a deeper

understanding of the population being studied (Ritchie & Levis, 2003).

### Instruments used

The measurement instruments were determined by the need to provide data from different perspectives to make the comparison between their results more meaningful. For the realization of our research, two measuring instruments were used, which are:

- **The depression measuring instrument** - is a standardized questionnaire by Seligman and has been officially used since 1975. This questionnaire has been prepared for all parents who have children with a neurodevelopmental diagnosis. In the first part of this questionnaire, the questions about the demographic data of the subjects are presented, while the second part of the questionnaire contains 17 statements of the closed type with 6 rating scales.
- **Anxiety measuring instrument** - is a standardized questionnaire by Charles. D. Spielberg, C.D. Edwards, J. Monturi & R. Lushene, in 1973. This questionnaire was prepared for all parents who have children with a neuro-developmental diagnoses. Through this questionnaire, we understand how disturbing each of the listed symptoms has been during the last 30 days, including today. This questionnaire has 20 closed-type questions with 4 rating scales.

### Reliability of measuring instruments

Cronbach’s Alpha method was used in this study. Cronbach’s Alpha coefficient, to assess the reliability of the instrument as a whole (Table 1).

The results show that the measuring instrument which measures depression satisfies the important condition for the application of the measuring instrument, respectively the reliability values for the Cronbach’s Alpha method  $\alpha=0.819$  and Gutman = 0.75 reach the reliability values  $\alpha=0.80$ . Whereas the measuring instrument which measures anxiety partially satisfies the important condition for the application of the measuring instrument, respectively the reliability values for the Cronbach’s Alpha method  $\alpha=0.75$  and Gutman = 0.75 reach the reliability values  $\alpha=0.80$ .

### Study sample

In this study, we included 300 parents of primary school children and preschool children with neuro-developmental

**Table 1:** The reliability coefficient of measuring instruments for depression and anxiety.

No. of events	Cronbach’s Alpha	Lambda					
		1	2	3	4	5	6
Depression 17	0.819	0.712	0.874	0.755	0.817		
Anxiety 20	0.755	0.714	0.761	0.821	0.727	0.845	0.912

disorders from several residential areas in Kosovo. The subjects that we sampled combine the characteristics for an adequate sample (gender of parents, age, place of residence, financial status, level of education, birth order and age of the child with neurodevelopmental disorders, number of primary school children and preschool children, diagnosis of primary school children and preschool children, they possess all the characteristics of the population where the research will be conducted, and the subjects have equal opportunities to be included in the sample (Table 2).

**Statistics processing**

All variables of interest were checked for normal distribution and then a decision was made as to the type of statistical analysis to be used. On the basis of these initial data, a more detailed analysis was carried out, looking for patterns and connections by comparing averages and exploring correlations. The data collected from the questionnaires were processed and analyzed with the SPSS-21 statistical program. The statistical analyzes undertaken in this study are descriptive and cognitive.

**RESULTS**

In this part of the study, we will present the results achieved during this research.

For the clearest and most logical presentation of the results from the statistical analyses, we will present the data in tabular form.

**Results related to differential analysis**

In the following, we will present the results related to the relationship between the level of depression and anxiety in parents with *primary school children and preschool children* diagnosed with neuro-developmental disorders, in which the request to test the difference between the respondents in the context of the research variables is included (Table 3).

The minimization of the score for each respondent in the general scale for the assessment of depression is 1, the maximum is 6, the Average A=4.8191, and SD =1.3898. Whereas, the minimization of the result for each respondent in the sub-scale for the assessment of anxiety is 1, maximum 4, average 4 A=3.8845 and SD= 0.8007.

**Results related to correlations**

In the following, we will present the results related to the relationship between the level of depression and anxiety in parents with primary school children and preschool children diagnosed with neuro-developmental disorders, as well as the relationship between socio-demographic characteristics: gender, calendar age, residence, and status. From the evidence that the correlative statistical analysis gives us, we have these results, which we will present below in the tabular form.

**Table 2:** Tabular presentation of the number of participants in the study, by gender, age, place of residence and their relative percentage

Parent	Gender		Age			Residence		
	Frequencies	%	Age	Frequencies	%	Residence	Frequencies	%
Mother	153	51.0%	20-25years	39	13.0	Village	81	27.0%
			26-30years	53	17.7			
			31-35years	53	17.7			
			36-40years	57	19.0			
			41-45years	29	9.7			
Father	147	49.0%	46-50years	28	9.3	City	219	73.0%
			51-55years	16	5.3			
			56-60years	16	5.3			
			61-64years	9	3.0			
Total	300	100%	Total	300	100%	Total	300	100%

**Table 3:** The average and the standard deviation of the overall level and variables.

Variables	Number (N)	Minimum	Maximum	Average (A)	Standard Deviation (SD)
Depression	300	1	6	4.8191	1.3898
Anxiety	300	1	4	3.8845	0.8007

**Table 4.1:** Correlation between depression and age

		Depression	Age
Depression	Pearson Correlation	1	0.673**
	Sig. (2-tailed)		000
	N	300	300
Age	Pearson Correlation	0.673**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 4.2:** Correlation between depression and gender

		Depression	Gender
Depression	Pearson Correlation	1	0.691**
	Sig. (2-tailed)		000
	N	300	300
Gender	Pearson Correlation	0.212**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 4.3.** Correlation between depression and residence

		Depression	Settlement
Depression	Pearson Correlation	1	0.364**
	Sig. (2-tailed)		000
	N	300	300
Settlement	Pearson Correlation	0.364**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 4.4.** Correlation between depression and financial status

		Depression	Financial status
Depression	Pearson Correlation	1	0.691**
	Sig. (2-tailed)		000
	N	300	300
Financial status	Pearson Correlation	0.691**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 4.5.** Correlation between depression and level of education

		Depression	Education level
Depression	Pearson Correlation	1	0.519**
	Sig. (2-tailed)		000
	N	300	300
Education level	Pearson Correlation	0.519**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 4:** Correlation between depression and sociodemographic characteristics: age, gender, place of residence, financial status, level of education in parents with primary school children and preschool children diagnosed with neuro-developmental disorders.

		Anxiety	Age
Anxiety	Pearson Correlation	1	0.221**
	Sig. (2-tailed)		000
	N	300	300
Age	Pearson Correlation	0.221**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 5:** Correlation between anxiety and socio-demographic characteristics: age, gender, place of residence, financial status, level of education in parents with primary school children and preschool children diagnosed with neuro-developmental disorders

**Table 6:** Tabular presentation of gender differences in the context of depression in parents

Variables	Gender	Number of cases (N)	Average (A)	Standard Deviation (SD)	Standard Error (SE)
Depression in parents	Mother	153	5.127	1.155	0.588
	Father	147	4.898	0.854	0.305

*Table 7. Tabular representation of gender differences in the context of anxiety in parents*

Variables	Gender	Number of cases (N)	Average (A)	Standard Deviation (DS)	Standard Error (SE)
Anxiety in parents	Mother	153	4.951	0.940	0.451
	Father	147	4.215	0.537	0.216

## DISCUSSION

From the results of the study and the analysis of the findings (see table 3), it appears that depression has a high positive cor-

		Anxiety	Gender
Anxiety	Pearson Correlation	1	0.148**
	Sig. (2-tailed)		000
	N	300	300
Gender	Pearson Correlation	0.148**	1
	Sig. (2-tailed)	000	
	N	300	300

*Table 5.1. Correlation between anxiety and age*

		Anxiety	Residence
Anxiety	Pearson Correlation	1	0.311**
	Sig. (2-tailed)		000
	N	300	300
Residence	Pearson Correlation	0.311**	1
	Sig. (2-tailed)	000	
	N	300	300

*Table 5.2. Correlation between anxiety and gender*

		Anxiety	Financial Status
Anxiety	Pearson Correlation	1	0.587**
	Sig. (2-tailed)		000
	N	300	300
Financial status	Pearson Correlation	0.587**	1
	Sig. (2-tailed)	000	
	N	300	300

**Table 5.4.** Correlation between anxiety and financial status

		Anxiety	Education level
Anxiety	Pearson Correlation	1	0.614**
	Sig. (2-tailed)		000
	N	300	300
Education level	Pearson Correlation	0.614**	1
	Sig. (2-tailed)	000	
	N	300	300

*Table 5.5. Correlation between anxiety and level of education*

relation with anxiety  $r=0.821$ ;  $p<0.01$  at the 0.01 significance level. As the level of depression in parents increases, so does the level of anxiety. From the evidence provided by correlative

statistical analysis, we can conclude that there is a high positive correlation between depression and anxiety.

Many authors have found associations between depression and anxiety in parents whose children have been diagnosed with neurodevelopmental disorders (Bornstein, 1993; Shuman et al., 1998). During the analysis of these studies, a significant connection between anxiety and depression in parents is found. This connection is confirmed by the studies of Stuart and Noyes (1999), where according to them, children's neurodevelopmental disorders are related to the manifestation of symptoms of anxiety disorders and depression in parents, the experience of psychological disturbances with physical symptoms. Somatizing behavior is best understood as a unique form of interpersonal behavior that results from neuropsychological disorders in their children (Stuart & Noyes, 1999). Depression affects different people in different ways. People suffering from depression may: be extremely angry and impatient, be engaged in negative thoughts, be bored, hateful, be under tension, lack security, blame themselves and feel guilty about things unnecessarily, smoke cigarettes more than usual, drink alcohol, and use narcotic substances; have decreased energy (have the feeling of losing energy); feeling worthless and useless; do not enjoy activities; lose sexual activity; feel bad spiritually most of the time; have suicidal thoughts; change in the way of eating; they eat too much and gain weight; or they don't eat well and lose weight; have difficulty concentrating or making decisions; they disconnect from others (isolate), instead of asking for help and support, etc. (Demyttenaere, 2008).

Women are more prone to depression than men. The highest risk for the female gender can be hormonal changes, which come as a result of puberty, the monthly cycle, menopause or pregnancy (Stoep & Weiss, 2003). Although the risk in men is lower, depression can go undiagnosed. This is due to the fact that many men mask their illnesses with alcoholic drinks, use drugs, abuse and violence. Suicide is a serious risk for males, 4 times more than females (Demyttenaere, 2008). From the results of the study, it appears that depression has a moderate positive correlation with the age of the parents  $r=0.673$ ;  $p<0.01$  at the significance level 0.01 (see table 4.1). From the evidence provided by the correlative statistical analysis, we can conclude that age can partially influence the decrease or increase in the degree of depression.

The results of the study also show us that depression has a weak positive correlation with the gender of the parents  $r=0.212$ ;  $p<0.01$  at the 0.01 significance level (see table 4.2). From the evidence provided by the correlative statistical analysis, we can conclude that there is a weak positive correlation between depression and the gender of the parents because the gender of the parents is not an important factor that can influence the increase or decrease in the level of depression.

The results also show that depression has a weak positive correlation with parents' residence  $r=0.364$ ;  $p<0.01$  at the 0.01

significance level (see table 4.3). From the evidence provided by the correlative statistical analysis, we can conclude that there is a weak positive correlation between depression and parents' residence for the reason that parents' residence is not an important factor that can influence the increase or decrease in the level of depression.

The results show that depression has a moderate positive correlation with the financial status of parents  $r=0.691$ ;  $p<0.01$  at the 0.01 significance level (see table 4.4). From the evidence provided by the correlative statistical analysis, we can conclude that there is a moderate positive correlation between depression and the financial status of the parents, it is moderate for the reason that financial status can partially influence the increase or decrease in the degree of depression.

Our results also show that depression has a moderate positive correlation with parents' education level  $r=0.519$ ;  $p<0.01$  at the 0.01 significance level (see table 4.5). From the evidence that the correlative statistical analysis gives us, we can conclude that there is a medium positive correlation between depression and the level of education of the parents, it is medium because the level of education can partially affect the increase or decrease of the level of depression.

In support of these data are many of the results of other studies, however, it is important to emphasize that the results regarding depression and sociodemographic characteristics must be interpreted carefully and certainly require repetition since some of these relationships do not persist after multiple comparisons.

A recent article (Olivari, Tagliabue, & Confalonieri, 2013), found authoritative parenting style positively associated with adaptive behaviors and psychological adjustment and negatively associated with maladaptive behaviors and psychological maladjustment. Other articles concurred or agreed with this statement, (Kawabata, Alink, Tseng, van IJendoorn, & Crick, 2011; Newman, Harrison, Dashiff, & Davies, 2008; Piko & Balázs, 2012), suggesting the role of data demographics that play in parents with children diagnosed with neurodevelopmental disorders (Tagliabue S., 2014).

From the results of our study regarding the connection between anxiety and socio-demographic characteristics: age, gender, residence, financial status, and level of education among parents with children diagnosed with neurodevelopmental disorders, it appears that anxiety has a weak positive correlation with the age of parents  $r=0.221$ ;  $p<0.01$  at the 0.01 significance level (see table 5.1).

Likewise, from the results of our study, we see that anxiety has a very weak positive correlation with the age of the parents  $r=0.148$ ;  $p<0.01$  at the 0.01 significance level (see table 5.2). From the evidence that the correlative statistical analysis gives us, we can conclude that there is a very weak positive connection between anxiety and the parents' age, it is a very weak connection for the reason that the parents' gender is

not an important factor that can influence the increasing or decreasing level of anxiety.

The results show that anxiety has a weak positive correlation with parents' residence  $r=0.311$ ;  $p<0.01$  at the 0.01 significance level (see table 5.3). From the evidence that the correlative statistical analysis gives us, we can conclude that there is a weak positive connection between anxiety and the residence of the parents, it is a weak connection for the reason that the residence of the parents is not an important factor that can influence the increasing or decreasing level of anxiety.

From our result we also see that anxiety has a moderate positive correlation with the financial status of parents  $r=0.587$ ;  $p<0.01$  at the 0.01 significance level (see table 5.4). From the evidence provided by the correlative statistical analysis, we can conclude that there is a medium positive connection between anxiety and the financial status of the parents, it is a medium connection because the financial status of the parents partially affects the increasing or decreasing level of anxiety.

Our results show that anxiety has a moderate positive correlation with parents' education level  $r=0.614$ ;  $p<0.01$  at the 0.01 significance level (see table 5.5). From the evidence that the correlative statistical analysis gives us, we can conclude that there is a medium positive connection between anxiety and the level of the parents' education, it is a medium connection because the level of the parents' education partially affects the increasing or decreasing level of anxiety.

There are many studies have been done regarding the connection between anxiety and sociodemographic data. In the study of Noel (2001), referring to the sociodemographic factors of the parents and the neuropsychological disorders of their children, it was observed that there was an association in the negative direction between the gender of the parents and anxiety, where fathers' result in higher levels of anxiety compared to mothers.

On the other hand, it is noticed in the study by Anderson (2000), a significant statistical association between the gender of the parent and parents' anxiety, where it results that fathers are more likely to show anxiety compared to mothers.

Russell (1998), cites data supporting age differences in parents' anxiety. There is also a negative connection between the age of children with neurodevelopmental disorders and parents' anxiety, where parental protection towards children seems to decrease as the child's age increases. In contrast to this, the positive direction of the connection between the age of the parent and the parenting style shows us a tendency toward the authoritarian style with increasing age, especially over 45 years old.

Parental education level affects parenting anxiety. Studies show that when the level of education increases, the democratic attitudes of parents towards children also increase, and the opposite happens when the level of education of parents is low, they experience more anxiety (Ari Bayhan & Artan, 1995).

From our results regarding the statistically significant gender differences between the level of depression and parents whose children have been diagnosed with neuro-developmental disorders, we see that mothers reach a higher average  $M=5.127$  than fathers  $M=4.898$ . The standard deviation for fathers is  $Ds=1.115$ , while for mothers  $Ds=0.854$ , and the average standard error for fathers is  $Se=0.588$  and for mothers  $Se=0.588$  (see table 6). So from these results, we understand that there are gender differences in the variables of depression in parents, but that there are no significant statistical differences.

The reason why mothers achieve a higher depression average than fathers is that mothers are more emotionally sensitive, they are also more isolated at home and spend more time with their children. These factors cause mothers to have higher average depression than fathers.

From our results regarding gender differences, statistically significant between the level of anxiety and parents whose children have been diagnosed with neuro-developmental disorders, we see that mothers reach a higher average  $A=4.951$  than fathers  $A=4.215$ . The standard deviation for fathers is  $Sd=0.940$ , while for mothers  $Sd=0.537$ , and the average standard error for fathers is  $Se=0.451$  and for mothers  $Se=0.216$  (see table 7). From these results, we can say that there are gender differences in the variables of anxiety in parents, but they do not have significant statistical differences.

The reason why mothers have higher average anxiety is the same as depression mothers are more emotionally sensitive, they are also more isolated at home and spend more time with their children. These factors cause mothers to have higher average depression than fathers.

## CONCLUSIONS

Do not rush to solve the problems that children face. Children have a natural need to face life's fears and challenges. If the parents will solve these problems for them, they will never feel strong and capable enough to solve them themselves and will later be dependent on external factors in solving the problems without taking their own initiative for this.

Children benefit from mistakes and testing skills and strengths through personal experiences. When they are over-instructed, they lose confidence in their abilities. Avoiding disappointments, concerns, and ejections failures make children vulnerable and unprepared for the future. Often fears and anxieties are not expressed but kept locked inside themselves, shying away from the judgment of others.

Identify your anxieties and worries. Anxiety is often a learned reaction and can be passed down from one generation to the next. If you are more aware of your own anxiety as a parent, you will take steps to deal with it rather than transferring it onto the child.

Teach the child relaxation techniques and autosuggestion. These techniques have quite good effects in relieving anxiety and moving to another state of mind where the person feels more positive and ready to deal with intrapersonal concerns.

## REFERENCES

- American Psychiatric Association, (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Anastasi, A., & Urbina, S. (1997). *Psychological testing* (7th ed). Upper Saddle River, NJ: Prentice Hall.
- Bartels, D. & Pizarro, D.A. (2011). The mismeasure of morals: Antisocial personality traits predict utilitarian responses to moral dilemmas. *Cognition*, 121, 154-161.
- Berger S., Kathleen (18 February 2011). *The Developing Person Through the Life Span* (8th ed.). Worth Publishers. pp. 273–278.
- Bernstein, Robert (20 February 2008). "Majority of Children Live With Two Biological Parents". Archived from the original on 20 April 2008. Retrieved 26 March 2009.
- Bögels & Brechman-Toussaint, 2006. Family issues in child anxiety: Attachment, family functioning, parental rearing and beliefs. *Clinical Psychology Review*, 26, 834– 856
- Bornstein MC, Zlotnik D., 2009. Parenting styles and their effects. In: Benson JB, Hiath MM, eds. *Social and Emotional Development in Infancy and Early Childhood*. San Diego, CA: Elsevier; 280 –292
- Bornstein, M (Ed.). (2002). *Handbook on parenting: children and parenting* (Vol. 1). Mahwah NJ: Lawrence Erlbaum Publishers.
- Brannan, A.M., Manteuffel, B.A., Holden, E.W., & Heflinger, C.A. (2006). Use of the Family Resource Scale in children's mental health: Reliability and validity in economically diverse samples. *Administration and Policy in Mental Health and Mental Health Services Research*, 33, 182-197.
- Chorpita, B. F., & Barlow, D. H. (1998). The development of anxiety: The role of control in the early environment. *Psychological Bulletin*, 124, 3–21.
- Constantinople, A. A. (1969). An Eriksonian measure of personality development in college student template: Unclear what this sentence is trying to convey. *Developmental Psychology*, 1357-372.
- Costello, E. J., Mustillo, S., Erkanli, A., Keeler, G., & Angold, A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of General Psychiatry*, 60(8), 837-844, (2003).
- Craske, M. G. (1999). *Anxiety disorders: Psychological approaches to theory and treatment*. Boulder, CO: Westview.
- Edwards, S. L., Rapee, R. M., & Kennedy, S. (2010). Prediction of anxiety symptoms in preschool-aged children: Examination of maternal and paternal perspectives. *Journal of Child Psychology and Psychiatry*, 51, 313–321.
- Eley, T. C. (2001). Contributions of behavioural genetics research: Quantifying genetic, shared environmental and nonshared environmental influences.
- Freud, Sigmund. *The Standard Edition of the Complete Psychological Works of Sigmund Freud*. Translated by James Strachey. 24 vols. London: Hogarth, 1953–74.
- Kawabata, Y., Alink, L.R.A., Tseng, W.L., Van IJzendoorn, M.H., & Crick, N.R. (2011). Maternal and paternal parenting styles associated with relational aggression in children and adolescents: A conceptual analysis and meta-analytic review. *Developmental Review*, 31, 240-278.
- Kendler et al, 1986. et al(1986) Symptoms of anxiety and depression in a volunteer twin population. *Archives of General Psychiatry*, 43, 213-221.
- Lanz, M., Scabini, E., Vermulst, A. A., & Gerris, J. R. M. (2001). Congruence on child rearing in families with early adolescent and middle adolescent children. *International Journal of Behavioral Development*, 25(2), 133–139.
- Larzelere, R. E. & Baumrind, D. (2010). Are spanking injunctions scientifically supported? *Law and contemporary problems*, 73, 57-74.
- Lavigne, Cromley T, Sprafkin J, Gadow KD., 2009. The Child and Adolescent Symptom Inventory-Progress Monitor: a brief Diagnostic and Statistical Manual of Mental Disorders, 4th edition-referenced parent-report scale for children and adolescents. *J Child Adolesc Psychopharmacol*. 2009 Jun;19(3):241-52. doi: 10.1089/cap.2008.052.
- Olivari, M. G., Tagliabue, S., & Confalonieri, E. (2013). Parenting Style and Dimensions Questionnaire: A Review of Reliability and Validity. *Marriage & Family Review*, 49, 465-490.
- Ollendick, T.H., Davis, T.E. & Muris, P. (2004). Treatment of specific phobia in children and adolescents. In P.M. Barrett & T.H. Ollendick (Eds.), *Handbook of interventions that work with children and adolescents: Prevention and treatment* (pp.273-299). New York.
- Phoebe S. Moore, Shannon E. Whaley, and Marian Sigman (2004), 'Interactions between mothers and children: Impacts of maternal and child anxiety', *Journal of Abnormal Psychology*, 113, No, 471-476.
- Rapee, R.M. (1997). Potential role of childrearing practices in the development of anxiety and depression. *Clinical Psychology Review*, 17, 47–67. Raskin et al, 1982. Panic and generalized anxiety disorder. *Archives of General Psychiatry*, 39, 687-689.
- Repetti, R.L. (1996). The effects of perceived daily social and academic failure experiences on school-age children's subsequent interactions with parents. *Child Development*, 67(4), 1467-1482.
- Rubin, K., & Stewart, S. (1996). Social withdrawal. *Child psychopathology* (pp. 277- 307).
- Sayal, K. Annotation: Pathways to care for children. *Journal of Child Psychology and Psychiatry*, 47:7, 649–659. (2006). Stoep, Weiss, e bp., 2003
- Shuman, L., Vandek, D., & Posner, J. (1998). Harsh, firm, and permissive parenting in low-income families. *Journal of Family Issues*, 19, 483-507.
- Stuart, S., & Noyes, R. (1999). Attachment and interpersonal communication in somatization disorder. *Psychosomatics*, 40, 34–43.
- Tamo A., Karaj Th., Rapti E., 2007. Practices of child-rearing in the remote north and suburban regions in Albania. Study Report Prepared by the Human Development Center, Tirana, Albania.
- U.S. Public Health Service, (2000). Report on the Surgeon General's Conference on Children's Mental Health: A National Action Agenda. Washington, DC:Fouche 2002:16
- Unger, J. B., Kipke, M. D., Simon, T. R., Montgomery, S. B., & Johnson, C. J. Homeless youths and young adults in los angeles: Prevalence of mental health problems and the relationship between mental health and substance abuse disorders. *American Journal of Community Psychology*, 25(3), 371-394, (1997).
- White, F., Hayes, B., & Livesey, D. (2005). *Developmental Psychology: From Infancy to Adulthood*. NSW: Pearson Education Australia.